

(19)



(11)

EP 1 983 046 A1

(12)

EUROPEAN PATENT APPLICATION
published in accordance with Art. 158(3) EPC

(43) Date of publication:

22.10.2008 Bulletin 2008/43

(51) Int Cl.:

C12N 15/00 ^(2006.01) **C12N 15/09** ^(2006.01)
C12Q 1/68 ^(2006.01)

(21) Application number: **07707949.9**

(86) International application number:

PCT/JP2007/051800

(22) Date of filing: **02.02.2007**

(87) International publication number:

WO 2007/088971 (09.08.2007 Gazette 2007/32)

(84) Designated Contracting States:

DE FR GB IT

(30) Priority: **03.02.2006 JP 2006027735**

(71) Applicants:

- **MessengerScape Co. Ltd.**
Shibuya-ku
Tokyo
151-0072 (JP)
- **ORIENTAL YEAST CO., LTD.**
Tokyo 174-8505 (JP)

- **MIKAMI, Yoji**
Tokyo 1510072 (JP)
- **KINUGASA, Masahiro**
Nagahama-shi, Shiga 5260804 (JP)
- **MORI, Kazuya**
Nagahama-shi, Shiga 5260804 (JP)
- **SUGIMOTO, Michiyo**
Nagahama-shi, Shiga 5260804 (JP)
- **UCHIDA, Koji**
Nagahama-shi, Shiga 5260804 (JP)

(72) Inventors:

- **SAITO, Toshiyuki**
Tokyo 1510072 (JP)

(74) Representative: **Graf von Stosch, Andreas et al**
Bosch Graf von Stosch Jehle
Patentanwalts-gesellschaft mbH
Prinzregentenstrasse 22
80538 München (DE)

(54) **GENE GROUP APPLICABLE TO CANCER PROGNOSTICATION**

(57) To provide a novel method for determining the risk of lymph node metastasis of breast cancer uses as an index the difference in the expression levels of marker genes between metastatic breast cancer tissue (or cell) and non-metastatic breast cancer tissue (or cell). The method involves (1) measuring the expression level of a gene having a base sequence in human metastatic breast cancer tissue (or cell), (2) measuring the expres-

sion level of the same gene in human non-metastatic breast cancer tissue (or cell), and (3) comparing the expression level of (1) with the expression level of (2) and determine the risk of lymph node metastasis of breast cancer based on the difference between the expression levels.

EP 1 983 046 A1

Description

Technical Field

5 **[0001]** The present invention relates to a novel method for determining the risk of lymph node metastasis of breast cancer. More specifically, the present invention relates to a method for determining the risk of lymph node metastasis of breast cancer that is based on comparison of the expression levels of marker genes having specific base sequences between metastatic breast cancer cells and non-metastatic breast cancer cells.

10 Background Art

[0002] In Japan, the number of breast cancer patients is growing rapidly. The cancer is the most prevalent of all cancers in women. Estrogen, a female hormone, is considered a risk factor of breast cancer: women who have been exposed to estrogen for a longer period of time due to early menarche, late menopause, late age at first birth or nulliparity are more likely to develop breast cancer. Western-style high-fat diet and obesity are also associated with this type of cancer since estrogen is primarily produced in fat tissue in postmenopausal women. The changing lifestyles of Japanese women, such as their active participation in society, also contribute to the increase in the incidence of breast cancer.

15 **[0003]** Breast cancer is generally divided into three classes: non-invasive carcinomas, invasive carcinomas and Paget's disease of the breast. Most of the incidences of breast cancer that form lumps are invasive. There are common and special types of invasive breast cancers. The common types include scirrhoma, papillotubular carcinoma and solid-tubular carcinoma. The special types include mucinous carcinoma.

[0004] Because no blood test is available to specifically detect breast cancer, the detection of early breast cancers relies primarily on palpation and X-ray imaging. However, these techniques, even when used in combination, fail to detect as much as 20% of the cancer. In addition, diagnosis by X-ray imaging often requires specialists. The cytodiagnosis conducted before and during the surgical procedures can only be done by a pathologist and is often difficult due to the shortage of experienced pathologists and varying standards of the diagnosis. Thus, no subjective and simple technique for the detection/diagnosis of early breast cancers has ever existed to bridge the gap between detection and diagnosis of the disease. The PET analysis, a new diagnostic technique that can detect tumor tissue 1mm or less in diameter, requires large-scale facilities and is therefore not readily used for the detection of breast cancer.

25 **[0005]** Recent studies have shown that cancers are caused by anomalies in genes. For example, techniques have been proposed that detect cancer cells by making use of the fact that certain genes are expressed at different levels in a cancer tissue and a normal tissue (Patent Literatures 1 and 2).

[Patent Literature 1] Japanese Patent Application Laid-Open (JP-A) No. 2003-284594

35 [Patent Literature 2] Japanese Patent Application Laid-Open (JP-A) No. 2003-284596

Disclosure of the Invention

[0006] Once lymph node-metastatic breast cancer has been removed by surgery, prognosis is predicted based on indices such as tumor size, nuclear pleomorphism of the removed cancer cells and of hormone receptor levels. Where necessary, adjuvant therapy is given to prevent metastasis to lymph nodes or the recurrence of cancer. The prediction of prognosis based on these presently available indices is not accurate enough, however, and more accurate indices for the prognosis of breast cancer patients are therefore needed to reduce the risk of recurrence and improve patients' quality of life by proper medication.

45 **[0007]** In view of the above-described problems, the present inventors have conducted extensive studies and observed that certain marker genes are expressed at different levels in metastatic breast cancer cells or tissues and in non-metastatic breast cancer cells or tissues. The present inventors found that these marker genes could be used to determine the risk of lymph node metastasis of breast cancer and ultimately devised the present invention. Accordingly, the present invention provides the following measures to address the above-described problems.

50 (1) A method for determining the risk of lymph node metastasis of breast cancer, including using as an index the difference in the expression level of a marker gene between a metastatic breast cancer tissue (or cell) and a non-metastatic breast cancer tissue (or cell).

(2) The method according to (1) above, wherein the expression level of the marker gene is determined by the amount of mRNA of the gene.

55 (3) The method according to (1) or (2) above, wherein the marker gene is at least one selected from the group consisting of genes having base sequences of GenBank accession Nos. NM000903, NM006804, NM033547, CR611676, NM177967, NM152558, NM178167, NM003752, AK131568, CR592336, NM178507, NM002862,

NM006913, NM005794, NM014164, NM000853 and a base sequence extending from 178882962bp to 178883181bp of chromosome 3, and homologs thereof.

(4) The method according to any one of (1) to (3) above, wherein the expression level of the marker gene in the metastatic breast cancer tissue (or cell) is equal to or higher than twice the expression level in the non-metastatic breast cancer tissue (cells), or equal to or lower than one-half the expression level in the non-metastatic breast cancer tissue.

[0008] The method of the present invention enables quick and simple determination of the risk of lymph node metastasis of breast cancer at the genetic level, thus providing an effective way to prevent metastasis of breast cancer.

Brief Description of the Drawings

[0009]

Fig. 1 is a diagram showing a comparison of the expression levels of a transcript (transcript 1) of a marker gene according to high-coverage gene expression profiling (HiCEP).

Fig. 2 is a diagram showing a comparison of the expression levels of a transcript (transcript 2) of another marker gene according to HiCEP.

Fig. 3 is a diagram showing a comparison of the expression levels of a transcript (transcript 3) of another marker gene according to HiCEP.

Fig. 4 is a diagram showing a comparison of the expression levels of a transcript (transcript 4) of another marker gene according to HiCEP.

Fig. 5 is a diagram showing a comparison of the expression levels of a transcript (transcript 5) of another marker gene according to HiCEP.

Fig. 6 is a diagram showing a comparison of the expression levels of a transcript (transcript 6) of another marker gene according to HiCEP.

Fig. 7 is a diagram showing a comparison of the expression levels of a transcript (transcript 7) of another marker gene according to HiCEP.

Fig. 8 is a diagram showing a comparison of the expression levels of a transcript (transcript 8) of another marker gene according to HiCEP.

Fig. 9 is a diagram showing a comparison of the expression levels of a transcript (transcript 9) of another marker gene according to HiCEP.

Fig. 10 is a diagram showing a comparison of the expression levels of a transcript (transcript 10) of another marker gene according to HiCEP.

Fig. 11 is a diagram showing a comparison of the expression levels of a transcript (transcript 11) of another marker gene according to HiCEP.

Fig. 12 is a diagram showing a comparison of the expression levels of a transcript (transcript 12) of another marker gene according to HiCEP.

Fig. 13 is a diagram showing a comparison of the expression levels of a transcript (transcript 13) of another marker gene according to HiCEP.

Fig. 14 is a diagram showing a comparison of the expression levels of a transcript (transcript 14) of another marker gene according to HiCEP.

Fig. 15 is a diagram showing a comparison of the expression levels of a transcript (transcript 15) of another marker gene according to HiCEP.

Fig. 16 is a diagram showing a comparison of the expression levels of a transcript (transcript 16) of another marker gene according to HiCEP.

Fig. 17 is a diagram showing a comparison of the expression levels of a transcript (transcript 17) of another marker gene according to HiCEP.

Best Mode for Carrying Out the Invention

[0010] The present invention concerns a method for determining the risk of lymph node metastasis of breast cancer that uses as an index of the risk of metastasis the difference in the expression levels of specific marker genes between metastatic breast cancer cells or tissues and non-metastatic breast cancer cells or tissues. As used herein, the term "marker gene" refers to a gene that enables the determination of the risk of metastasis of breast cancer cells by comparing its expression levels between metastatic breast cancer cells or tissues and non-metastatic breast cancer cells or tissues.

[0011] The present invention also concerns a method for determining the risk of lymph node metastasis of breast cancer in which the expression levels of the marker genes are determined by the amounts of mRNA of the marker genes.

More specifically, the present invention concerns a method for determining the risk of lymph node metastasis of breast cancer that involves extracting total RNA from cells obtained from metastatic and non-metastatic breast cancer tissues, and comparing the amounts of mRNA transcripts transcribed from the marker genes. Different techniques for gene expression analysis can be used to determine the amounts of mRNA of genes of interest, including comprehensive transcriptome analysis (high-coverage gene expression profiling, HiCEP), DNA microarrays, quantitative RT-PCR and northern hybridization. Gene expression analysis techniques that can determine the amounts of mRNA without extracting total RNA from cells, such as *in situ* hybridization, may also be used in the present invention. The above-described techniques may be used in combination to improve the accuracy of detection. The translated products of the genes of the present invention may also be quantified by, for example, determining the amounts of proteins coded by the genes. Proteins of interest can be quantified by using techniques such as immunological assays using antibodies specific for the proteins (such as ELISA, western blotting and RIA), two-dimensional electrophoresis and high-performance liquid chromatography (HPLC). Antibodies specific for the proteins coded by the genes of the present invention can be prepared by common techniques using the proteins coded by the genes as antigens.

[0012] HiCEP is one of the transcriptome analysis techniques and is characterized by its comprehensiveness and high sensitivity. The following is a brief overview of the technique (See *Nucleic Acids Res.*, 2003, Vol. 31, No.16 e94 for more details): Using common techniques, total RNA is extracted and purified from tissue or cell samples. Double-stranded cDNA is synthesized from the total RNA using 5'-biotinylated oligo(dT) primers. The cDNA is then digested with a restriction enzyme MspI. Poly(A)-containing fragments are collected by avidin beads and 3'-adaptor is ligated to the MspI-digested ends of the collected fragments. The fragments are then digested with a restriction enzyme MseI and 3'-adaptor is ligated to the MseI-digested ends. PCR primers are constructed by adding all possible combinations of two selected bases to the same adaptor sequences as those ligated to 5' and 3' ends (16 5'-end primers and 16 3'-end primers with 5'-end primers fluorescent-labeled). Using these primers, 256 different quantitative PCRs are performed. The PCR products obtained for each primer pair are loaded on a fragment analyzer to obtain 256 electrophoresis profiles (gene expression profiles), each containing multiple fluorescence peaks, for a sample. The expression levels of transcripts can then be compared by comparing the fluorescence peaks among different samples.

[0013] The marker gene for use in the present invention may be any gene that is expressed at significantly different levels between metastatic breast cancer cells or tissues and non-metastatic breast cancer cells or tissues. For example, the marker gene may be at least one selected from the group consisting of genes having base sequences of GenBank accession Nos. NM000903, NM006804, NM033547, CR611676, NM177967, NM152558, NM178167, NM003752, AK131568, CR592336, NM178507, NM002862, NM006913, NM005794, NM014164 and NM000853 and a base sequence extending from 178882962bp to 178883181bp of chromosome 3, and homologs thereof.

[0014] Data stored in the GenBank database may contain the same gene registered by different researchers, at different times, in different fields and under different names or gene polymorphisms or splicing variants of the same gene registered as novel genes. Thus, different base sequences that can be considered to be originated from a single gene may be registered with different accession numbers. These base sequences are collectively referred to as "homologs." The term is used in the same context throughout this specification.

[0015] One characteristic feature of the method of the present invention for determining the risk of lymph node metastasis of breast cancer is the use of marker genes that are expressed at significantly different expression levels between metastatic breast cancer cells or tissues and non-metastatic breast cancer cells or tissues. The term "expression level" may refer to either the amount of mRNA transcribed from a marker gene or the amount of a protein translated from mRNA. With regard to the difference in the expression level of a marker gene between metastatic breast cancer cells or tissues and non-metastatic breast cancer cells or tissues, the ratio of the expression level of a marker gene in non-metastatic breast cancer cells or tissues to the expression level of the same gene in metastatic breast cancer cells or tissues is preferably in the range of 1.5 or higher or 2/3 or lower, and more preferably in the range of 2 or higher or 1/2 or lower. A marker gene does not serve as an accurate index of the risk of lymph node metastasis of breast cancer and is therefore not desirable when the ratio of its expression level in non-metastatic breast cancer cells or tissues to that in metastatic breast cancer cells or tissues is outside the above-described range.

[0016] While the method of the present invention can be applied to any type of breast cancer, including breast ductal carcinomas (such as papillotubular carcinoma, solid-tubular carcinoma and scirrhoma), lobular carcinomas, special-type carcinomas (such as mucinous carcinoma, medullary carcinoma and tubular carcinoma) and Paget's disease of the breast, it is preferably applied to scirrhoma, lobular carcinomas or solid-tubular carcinoma.

Example

[0017] The present invention will now be described with reference to Example, which is not intended to limit the scope of the invention in any way.

[0018] In this Example, the expression levels (RNA transcription levels) of different genes are compared between human metastatic breast cancer tissue and non-metastatic tissue using one of the gene expression analysis techniques

EP 1 983 046 A1

known as high-coverage gene expression profiling technique (HiCEP), a known comprehensive, highly sensitive technique for transcriptome analysis (Nucleic Acids Res., 2003, vol.31(16), e94).

[0019] The breast cancer tissues used in Example were shown in Table 1. The tissues were collected from five stage II breast cancer patients (commercial products, all Caucasian, primary tumor, lymph node metastasis (2), no lymph node metastasis (3), all had stage II cancer based on TNM classification).

Table 1

Samples	Metastasis	Tumor size	Age
#A	+	12cm	57
#B	+	2cm x 1.5cm x 1.5cm	69
#C	-	2.5cm	50
#D	-	4cm x 2cm x 1.7cm	61
#E	-	6cm x 5.5cm x 4.5cm	68

[0020] Total RNA was extracted from the samples by a common kit technique using RNeasy kit (Qiagen). 0.1µg of total RNA from each sample was used as template and reverse-transcribed using Super Script First Strand Synthesis System for RT-PCR (Invitrogen). The reverse transcript was incubated with DNA polymerase I (80 units), RNAase H (4 units, Invitrogen) and E. coli DNA ligase (40 units, Invitrogen) at 16°C for 2 hours. The resulting double-stranded DNA was incubated with restriction enzymes Mse I (40 units, New England Biolabs) and Msp I (50 units, TaKaRa Bio) at 37°C for 4 hours. Adaptor sequences were ligated to the ends of the resulting DNA fragments. Selective PCRs were performed using the adaptor-ligated DNA fragments as templates. The amplified products were analyzed by capillary electrophoresis. The waveform data were used to determine gene expression levels, compare the gene expression levels among the samples, and classify the genes into different expression patterns to obtain data for clustering (expression variation peaks).

[0021] The results of the analysis shown in Table 2 and Figs. 1 through 17 demonstrate that the difference in the fluorescence peak intensity between samples obtained from patients with lymph node metastasis and samples obtained from patients with no metastasis was significant for each of the 17 marker gene transcripts. In this analysis, each sample was assayed in two replicates and the resulting fluorescence peaks were overlapped. Arrows indicate the peaks for the marker gene transcripts.

[0022] Specifically, Transcripts 1 through 11 (as numbered in Table 1) each show a significant fluorescence peak in each of the metastasis samples but show no expression peak or, if any, a peak intensity that is half or less of the peak intensity of the metastasis samples in each of the non-metastasis samples. Conversely, Transcripts 12 through 17 each show a significant fluorescence peak in each of the non-metastasis samples but show no expression peak or, if any, a peak intensity that is half or less of the peak intensity of the non-metastasis samples in each of the metastasis samples. These observations demonstrate that each of the 17 genes can serve as an index of the risk of breast cancer metastasis that allows the determination of the risk of metastasis based on their expression levels.

EP 1 983 046 A1

Table 2. Transcripts as markers for breast cancer metastasis

Transcripts	Sequences	GenBank Accession No.	Annotation	Characteristics	
5	#1	Transcript sequence containing a base sequence from 68302490bp to 68317861bp of (-) strand of chromosome 16	NM_000903	NAD(P)H menadione oxidoreductase 1	Experssion enhanced in metastatic breast cancer
10	#2	Transcript sequence containing a base sequence from 178882962bp to 178883181bp of (+) strand	N/A	N/A	
15	#3	Transcript sequence containing a base sequence from 35050592bp to 35050643bp of (+) strand of chromosome 17	NM_006804	steroidogenic acute regulatory protein related	
20	#4	Transcript sequence containing a base sequence from 77267542bp to 77272569bp of (-) strand of chromosome 11	NM_033547	Homo sapiens hypothetical gene MGC16733 similar to CG12113 (MGC16733), mRNA.	
25	#5	Transcript sequence containing a base sequence from 176665540bp to 176666255bp of (+) strand of chromosome 5	CR611676	Similar to Pxl9-like protein (25 kDa protein of relevant evolutionary and lymphoid interest) (PRELI) (CGI-106) (SBB12)	
30	#6	Transcript sequence containing a base sequence from 98835662bp to 98835862bp of (+) strand of chromosome 13	NM_177967	Phosphoglycerate dehydrogenase like 1	
35	#7	Transcript sequence containing a base sequence from 2426581bp to 2426860bp of (+) strand of chromosome 7	NM_152558	IQ motif containing E (IQCE)	
40	#8	Transcript sequence containing a base sequence from 1987788 bp to 1987865 bp of (-) strand of chromosome 16	NM_178167	Zinc finger protein 598	
45	#9	Transcript sequence containing a base sequence from 228320033 bp to 28320077 bp of (-) strand of chromosome 16	NM_003752	Eukaryotic translation initiation factor 3, subunit 8, 110kDa	
50	#10	Transcript sequence containing a base sequence from 35135544 bp to 35135831 bp of (+) strand of chromosome 17	AK131568	V-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian)	
55					

EP 1 983 046 A1

(continued)

Transcripts	Sequences	GenBank Accession No.	Annotation	Characteristics
5 #11	Transcript sequence containing a base sequence from 35126382 bp to 35127393 bp of (+) strand of chromosome 17	CR592336	V-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian)	
10 #12	Transcript sequence containing a base sequence from 119605680 bp to 119605847 bp of (+) strand of chromosome 11	NM_178507	NS5ATP13TP2 protein	Expression decreased in metastatic breast cancer
15 #13	Transcript sequence containing a base sequence from 25226174 bp to 25226624 bp of (+) strand of chromosome 20	NM_002862	Phosphorylase, glycogen; brain	
20 #14	Transcript sequence containing a base sequence from 32256007 bp to 32256297 bp of (+) strand of chromosome 6	NM_006913	Ring finger protein 5	
25 #15	Transcript sequence containing a base sequence from 23183541 bp to 23184510 bp of (-) strand of chromosome 14	NM_005794	Dehydrogenase/reductase (SDR family) member 2	
30 #16	Transcript sequence containing a base sequence from 40352503 bp to 40352595 bp of (+) strand of chromosome 19	NM_014164	FXD domain containing ion transport regulator 5	
35 #17	Transcript sequence containing a base sequence from 22700873 bp to 22700983 bp of (+) strand of chromosome 22	NM_000853	Glutathione S-transferase theta 1	

40 Industrial Applicability

[0023] Genes according to the present invention enable the highly sensitive and subjective, yet simple and quick determination of lymph node metastasis of breast cancer, a task that has never been achieved by any of the conventional techniques. The genes of the present invention therefore serve as markers for the prognosis of breast cancer.

45

Claims

- 50 1. A method for determining the risk of lymph node metastasis of breast cancer, comprising: using as an index the difference in the expression level of a marker gene between a metastatic breast cancer tissue (or cell) and a non-metastatic breast cancer tissue (or cell).
- 55 2. The method according to claim 1, wherein the expression level of the marker gene is determined by the amount of mRNA of the gene.
3. The method according to claim 1 or 2, wherein the marker gene is at least one selected from the group consisting of genes having base sequences of GenBank accession Nos. NM000903, NM006804, NM033547, CR611676, NM177967, NM152558, NM178167, NM003752, AK131568, CR592336, NM178507, NM002862, NM006913,

EP 1 983 046 A1

NM005794, NM014164, NM000853 and a base sequence extending from 178882962bp to 178883181bp of chromosome 3, and homologs thereof.

- 5 **4.** The method according to any of claims 1 to 3, wherein the expression level of the marker gene in the metastatic breast cancer tissue (or cell) is equal to or higher than twice the expression level in the non-metastatic breast cancer tissue (cells), or equal to or lower than one-half the expression level in the non-metastatic breast cancer tissue.

10

15

20

25

30

35

40

45

50

55

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16





FIG. 17



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2007/051800

A. CLASSIFICATION OF SUBJECT MATTER C12N15/00(2006.01) i, C12N15/09(2006.01) i, C12Q1/68(2006.01) i		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) C12N15/00, C12N15/09, C12Q1/68		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Jitsuyo Shinan Koho 1922-1996 Jitsuyo Shinan Toroku Koho 1996-2007 Kokai Jitsuyo Shinan Koho 1971-2007 Toroku Jitsuyo Shinan Koho 1994-2007		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) PubMed, JMEDPlus (JDream2), JSTPlus (JDream2)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Yoji MIKAMI et al., "Hokatsuteki Kokando Tensha Sanbutsu Profile Kaiseki Gijutsu (HiCEP Gijutsu) o Shiyo shita Gan Maker Tansaku", The Japanese Journal of Clinical Pathology, Vol.53 Hosatsu, 2005, p.363	1-4
X	JP 2003-325192 A (Ortho-Clinical Diagnostics, Inc.), 18 November, 2003 (18.11.03), All pages (particularly, example 3) & EP 1367138 A2 & US 2003/0190656 A1	1, 2, 4
X	JP 2004-151003 A (Yuko NOGI), 27 May, 2004 (27.05.04), All pages (particularly, Claims 3, 7; referential example 2; Fig. 1) (Family: none)	1, 2, 4
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:		
"A"	document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed	
Date of the actual completion of the international search 20 February, 2007 (20.02.07)		Date of mailing of the international search report 27 February, 2007 (27.02.07)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (April 2005)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2007/051800

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	FUKUMURA, R., et al., A sensitive transcriptome analysis method that can detect unknown transcripts, Nucleic Acids Res., Vol.31, No.16, 2003, e94, [on line]. Retrieved from the Internet:<URL:http://www.pubmedcentral.nih.gov/picrender.fcgi?artid=169986&blobtype=pdf>.	1-4
A	JP 2003-043046 A (The Chemo-Sero-Therapeutic Research Institute), 13 February, 2003 (13.02.03), All pages (Family: none)	1-4
A	WO 2001/069260 A2 (RESEARCH FOUNDATION FOR MENTALHYGIENE, INC.), 20 September, 2001 (20.09.01), All pages & JP 2003-527607 A & EP 1269193 A1 & US 2003/0211554 A1	1-4
P,X	Yoji MIKAMI et al., "HiCEP ni yoru Nyugan Ten'isei Kanren Idenshi no Morateki Hatsugen Kaiseki", The Japan Radiation Research Society Taikai Koen Yoshishu, Vol.49, 2006.09.06, p.107	1-4
P,X	Yoji MIKAMI et al., "Ten'isei Nyugan to erbB2 Variant Idenshi Hatsugen no Kankei", Nyugan Kiso Kenkyu, Vol.15, 2006.04.01, p.7-11	1-4

Form PCT/ISA/210 (continuation of second sheet) (April 2005)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/JP2007/051800

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

The invention of claim 3 relates to a method for determining the possibility of lymph node metastasis of breast cancer by selecting one gene from 17 genes represented by GenBank Accession No. NM000903 and the like. However, the method for determining breast cancer metastasis (lymph node metastasis) using a specific gene as a marker is known as described in the documents 1-3 cited in the International Search Report. Therefore, a special technical feature, i.e., a technical feature (PCT Rule 13.2) that defines a contribution which each of the inventions, considered as a whole, makes over the prior art, does not exist in the method for determining the possibility of lymph node metastasis of breast cancer by selecting any gene from 17 (continued to extra sheet)

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee..
 The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2007/051800

Continuation of Box No.III of continuation of first sheet (2)

genes described in claim 3.

Thus, claim 3 includes 17 different inventions.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- JP 2003284594 A [0005]
- JP 2003284596 A [0005]

Non-patent literature cited in the description

- *Nucleic Acids Res.*, 2003, vol. 31 (16), e94 [0012]
[0018]